#KnowWhereToGo for children's cold and flu symptoms

Health Links - Info Santé

Call 204-788-8200 1-888-315-9257 (toll-free)

This free service is available 24/7 with nurses who can assess and provide advice on where to go.

Walk-In Connected Care Clinic

(After Hours & Weekends Winnipeg)

McGregor Walk-in and Access Fort Garry

When to get tested for COVID?



Doctor/Clinic/Primary Care (Weekdays)

During respiratory virus season, your child may experience common cold/flu symptoms.

- Fever or chills
- Cough
- Runny nose
- Sore throat

Critical Injury or Life-Threatening Condition?

Call 911 or go to an emergency department.



Doctor/Clinic/Primary Care Weekdays

After Hours and Weekends (Winnipeg)

Emergency Department / 911 24/7

During respiratory virus season, your child may experience common cold/flu symptoms. As long as symptoms are not emergent or urgent, they may be assessed by a primary care provider or clinic.

Mild symptoms may include:

- Fever or chills
- Nasal congestion or runny nose
- Cough
- Sore throat
- Fatique
- Headache
- Muscle aches or body aches
- Nausea or vomiting
- Diarrhea
- Poor appetite
- New loss of taste or smell
- Belly pain

McGregor Walk-in Connected Care Clinic 363 McGregor St

204-940-1963

Call ahead to confirm hours or check wait times

Monday - Friday: 12:00 p.m. - 7:30 p.m.

Saturday, Sunday and Holidays: 9:00 a.m. - 4:30 p.m.

Access Fort Gary Walk-in Connected Care Clinic 135 Plaza Dr

204-940-7100

Call ahead to confirm hours or check wait times

Monday – Friday: 9:00 a.m. - 6:30 p.m.

Saturday: 9:00 a.m. - 3:30 p.m.

Call 911 or go to your closest Emergency Department if your child has any of the following symptoms:

- Difficulty breathing
- Periods of holding breath / not breathing / slowed breathing
- Increased breathing (more than 60 breaths per minute)
- Excessively tired or difficult to wake up
- Sudden confusion
- Change in colour of nail beds, hands, lips, or face
- Inconsolable infants crying that won't stop or is painful sounding
- Not peeing
- Poor feeding refusing or unable to eat or drink
- Infant younger than 3 months with a fever