

JLCD-E-**AUTHORIZATION FOR ADMINISTRATION OF RELIEVE** MEDICATION & ASTHMA STANDARD HEALTH CARE PLAN (SHC



-11	
ΞR	
P)	
R١	

XI	Pembina Trails
	School Division Accomplish Anything
	Accomplish Anything

School Name:		School Year:
Student Information		
Name:	Birth	date:///
Address:		Year Month Day
MHSC# (6 digit):	PHIN	I # (9 digit):
Parent/Guardian/Caregiver Information		
Parent/Guardian/Caregiver:	Daytime phone(s)	
		Daytime phone(s)
		Daytime phone(s)
Medical Information		
Name	Dose	Medication device
☐ Salbutamol (e.g. Ventolin*, Airomir)	☐ 1 puff	☐ Metered dose inhaler (MDI)
☐ Symbicort*	☐ 2 puffs	☐ MDI & spacer device with mouthpiece ◀
□ Other	☐ 1 or 2 puffs	☐ MDI & spacer device with mask 🎉
School to contact URIS nurse if Parent/ Guardian/Caregiver selected "other".		☐ Turbuhaler
		□ Other
Name of prescribing physician:		
Trigger(s) for asthma (if known):		
Location of reliever medication:		all carry urgently required medication on their
change in medication.The pharmacy label must be on	the medication dev	annually with student registration or upon a rice. eplacing expired medication as well as the
I hereby request and authorize the schoutlined in the attached Asthma Stand Parent/Guardian/Caregiver Signature	ard Health Care Pla	e medication named above to my child as n.
School Administrator Signature:	Date	



Asthma Standard Health Care Plan

The Asthma SHCP is based on the clinical practice guidelines developed by the Unified Referral and Intake System (URIS) in consultation with Children's Allergy and Asthma Education Centre (CAAEC) at Health Sciences Centre. These clinical practice guidelines are available on the URIS website.

<u>Unified Referral and Intake System (URIS) | Manitoba Education and Early Childhood Learning (gov.mb.ca)</u>

IF YOU SEETHIS:	DOTHIS:	
 • Coughing • Wheezing • Chest tightness • Shortness of breath • Increase in rate of breathing while at rest 	 Remove the child from triggers of asthma. Have the child sit down. Ensure the child takes reliever medication (usually blue cap on bottom). Encourage slow deep breathing. Monitor the child for improvement of asthma symptoms. If reliever medication has been given and asthma symptoms do not improve in 5-10 minutes, contact the Parent/Guardian/Caregiver. Reliever medication can be repeated once at this time. If the child is not well enough to remain at the school, the Parent/Guardian/Caregiver should come and pick them up. If any of the emergency situations occur (see list below), call 911/EMS. 	
 Skin pulling in under the ribs Skin being sucked in at the ribs or throat Greyish/bluish colour in lips and nail beds Inability to speak in full sentences Shoulders held high, tight neck muscles Cannot stop coughing Difficulty walking 	 Activate 911/EMS. Delegate this task to another person. Do not leave the child alone. Continue to give reliever medication as prescribed every five minutes. Notify the Parent/Guardian/Caregiver. Stay with the child until EMS personnel arrive. 	